

-1-

## IYGB-MPI PAPER K - QUESTION 1

REWRITE IN INDEX NOTATION

$$y = \frac{8}{x} + 3\sqrt{x}$$

$$y = 8x^{-1} + 3x^{\frac{1}{2}}$$

$$\frac{dy}{dx} = -8x^{-2} + \frac{3}{2}x^{-\frac{1}{2}}$$

$$\frac{dy}{dx} = -\frac{8}{x^2} + \frac{3}{2\sqrt{x}}$$

SUBSTITUTING x=4

$$\frac{dy}{dx} \Big|_{x=4} = -\frac{8}{4^2} + \frac{3}{2\sqrt{4}}$$

$$= -\frac{8}{16} + \frac{3}{4}$$

$$= -\frac{1}{2} + \frac{3}{4}$$

$$= \frac{1}{4}$$



+-

## IYGB-MPI PAPER K - QUESTION 2

### a) COMPLETING THE SQUARE

$$f(x) = 4x^2 + 4x - 1$$

$$f(x) = 4\left[x^2 + x - \frac{1}{4}\right]$$

$$f(x) = 4\left[\left(x + \frac{1}{2}\right)^2 - \frac{1}{4} - \frac{1}{4}\right]$$

$$f(x) = 4\left(x + \frac{1}{2}\right)^2 - 1 - 1$$

$$\underline{f(x) = 4\left(x + \frac{1}{2}\right)^2 - 2}$$

ALTERNATIVE

$$\begin{cases} f(x) = 4x^2 + 4x - 1 \\ f(x) = (4x^2 + 4x + 1) - 2 \\ f(x) = (2x + 1)^2 - 2 \end{cases}$$

### b) SOLVING THE EQUATION USING PART (a)

$$f(x) = 0$$

$$4x^2 + 4x - 1 = 0$$

$$4\left(x + \frac{1}{2}\right)^2 - 2 = 0$$

$$4\left(x + \frac{1}{2}\right)^2 = 2$$

$$\left(x + \frac{1}{2}\right)^2 = \frac{1}{2}$$

$$x + \frac{1}{2} = \pm \sqrt{\frac{1}{2}} = \pm \frac{1}{\sqrt{2}}$$

$$x + \frac{1}{2} = \pm \frac{\sqrt{2}}{2}$$

$$x = -\frac{1}{2} \pm \frac{\sqrt{2}}{2}$$

ALTERNATIVE

$$\begin{cases} f(x) = 0 \\ (2x + 1)^2 - 2 = 0 \\ (2x + 1)^2 = 2 \\ 2x + 1 = \pm \sqrt{2} \\ 2x = -1 \pm \sqrt{2} \\ x = -\frac{1}{2} \pm \frac{\sqrt{2}}{2} \end{cases}$$

-1-

## IYGB - MPI PAPER K - QUESTION 3

PROCEED AS FOLLOWS

$$f(x) = x^{\frac{3}{2}} - 8x^{-\frac{1}{2}}$$

$$f(3) = 3^{\frac{3}{2}} - 8 \times 3^{-\frac{1}{2}} = (\sqrt{3})^3 - \frac{8}{\sqrt{3}} = 3\sqrt{3} - \frac{8}{\sqrt{3}}$$

$$= 3\sqrt{3} - \frac{8\sqrt{3}}{\sqrt{3}\sqrt{3}} = 3\sqrt{3} - \frac{8\sqrt{3}}{3} = 3\sqrt{3} - \frac{8}{3}\sqrt{3}$$

$$= \cancel{\frac{1}{3}\sqrt{3}}$$

1. E  $k = \frac{1}{3}$


-|-

## IYGB - MPI PAPER K - QUESTION 4

a) EXPANDING USING PASCAL'S TRIANGLE OR CALCULATOR

$$(2x+4)^3 = 1(2x)^3(4)^0 + 3(2x)^2(4)^1 + 3(2x)^1(4)^2 + 1(2x)^0(4)^3$$

$$\underline{(2x+4)^3 = 8x^3 + 48x^2 + 96x + 64}$$



b) USING PART (a)

$$(2x-1)(2x+4)^3 = (2x-1)(8x^3 + 48x^2 + 96x + 64)$$

$$= 16x^4 + 96x^3 + 192x^2 + 128x - 8x^3 - 48x^2 - 96x - 64$$

$$\underline{16x^4 + 88x^3 + 144x^2 + 32x - 64}$$

-1-


## IYGB - MPI PAPER K - QUESTION 5

a) START WITH + DIAGRAM

$$\vec{OA} = \underline{i} + 4\underline{j}$$

$$\vec{OB} = 5\underline{i} + 5\underline{j}$$

$$\vec{CB} = -\underline{i} + 6\underline{j}$$



FIND  $\vec{AB}$ , followed by its length

$$\vec{AB} = \vec{AO} + \vec{OB} = -(\underline{i} + 4\underline{j}) + (5\underline{i} + 5\underline{j}) = 4\underline{i} + \underline{j}$$

$$|\vec{AB}| = |4\underline{i} + \underline{j}| = \sqrt{4^2 + 1^2} = \underline{\sqrt{17}}$$

b) NEXT FIND THE VECTOR  $\vec{AC}$

$$\vec{AC} = \vec{AB} + \vec{BC} = \vec{AB} - \vec{CB} = (4\underline{i} + \underline{j}) - (-\underline{i} + 6\underline{j}) = 5\underline{i} - 5\underline{j}$$

NEXT THE MODULI OF  $\vec{AC}$

$$|\vec{AC}| = |5\underline{i} - 5\underline{j}| = \sqrt{5^2 + (-5)^2} = \sqrt{25 + 25} = \underline{\sqrt{50}}$$

c) FINALLY THE LENGTH OF CB

$$|\vec{CB}| = |-\underline{i} + 6\underline{j}| = \sqrt{(-1)^2 + 6^2} = \sqrt{1 + 36} = \sqrt{37}$$

BY THE COSINE RULE

$$\Rightarrow |\vec{AC}|^2 = |\vec{AB}|^2 + |\vec{CB}|^2 - 2|\vec{AB}||\vec{CB}|\cos\theta$$

$$\Rightarrow (\sqrt{50})^2 = (\sqrt{17})^2 + (\sqrt{37})^2 - 2\sqrt{17}\sqrt{37}\cos\theta$$

$$\Rightarrow 50 = 17 + 37 - 2\sqrt{629}\cos\theta$$

$$\Rightarrow 2\sqrt{629}\cos\theta = 4$$

$$\Rightarrow \cos\theta = 0.079745\dots$$

$$\therefore \underline{\theta = 85.4^\circ}$$

-1-

## IYGB-MPI PAPER K - QUESTION 6

a) LET  $f(x) = x^3 + x^2 - 10x + 8$

looking for factors, trying  $\pm 1, \pm 2, \pm 4, \pm 8$

$$f(1) = 1 + 1 - 10 + 8 = 0$$

$\therefore (x-1)$  is a factor of  $f(x)$

BY LONG DIVISION OR MANIPULATIONS


$$\begin{aligned} f(x) &= x^3 + x^2 - 10x + 8 = x^2(x-1) + 2x(x-1) - 8(x-1) \\ &= (x-1)(x^2 + 2x - 8) \\ &= (x-1)(x-2)(x+4) \end{aligned}$$

b) COLLECTING ALL INFORMATION

$$+x^3 \Rightarrow \curvearrowleft$$

$$x=0, y=8 \Rightarrow (0, 8)$$

$$y=0, x=\begin{matrix} 1 \\ 2 \\ -4 \end{matrix} \Rightarrow \begin{matrix} (1, 0) \\ (2, 0) \\ (-4, 0) \end{matrix}$$




- 1 -

## IYGB-MPI PAPER K - QUESTION 7

Process as follows

$$\begin{aligned}\text{GRADIENT} &= \frac{4\ln 5 - 2\ln 5}{2 - 0} \\ &= \frac{2\ln 5}{2} \\ &= \ln 5\end{aligned}$$



THE EQUATION OF THE STRAIGHT LINE IS

$$\Rightarrow \ln y - 2\ln 5 = (\ln 5)(t - 0)$$

$$\Rightarrow \ln y - 2\ln 5 = t\ln 5$$

$$\Rightarrow \ln y = t\ln 5 + 2\ln 5$$

$$\Rightarrow \ln y = \ln 5^t + \ln 5^2$$

$$\Rightarrow \ln y = \ln(5^t \times 5^2)$$

$$\Rightarrow \ln y = \ln(5^{t+2})$$

$$\Rightarrow y = 5^{t+2}$$

- -

## IYGB-MPI PAPER K - QUESTION 8


SOLVING THE FIRST INEQUALITY

$$\begin{aligned}\Rightarrow (x+2)(x+4) &> 10x + 7 \\ \Rightarrow x^2 + 6x + 8 &> 10x + 7 \\ \Rightarrow x^2 - 4x + 1 &> 0\end{aligned}$$

THIS DOES NOT FACTORIZE NICELY - OBTAIN CRITICAL VALUES BY  
COMPLETING THE SQUARE IN THE CORRESPONDING EQUATION - OR  
USE QUADRATIC FORMULA

$$\begin{aligned}\Rightarrow "x^2 - 4x + 1 = 0" \\ \Rightarrow (x-2)^2 - 4 + 1 = 0 \\ \Rightarrow (x-2)^2 = 3 \\ \Rightarrow x-2 = \pm\sqrt{3} \\ \Rightarrow x = 2 \pm \sqrt{3}\end{aligned}$$

LOOKING AT THE DIAGRAM



SOLVING THE SECOND INEQUALITY

$$\begin{aligned}\Rightarrow 3x\sqrt{3} < 2 + \frac{2(2x-1)}{\sqrt{3}} \\ \Rightarrow 3x < 2\sqrt{3} + 2(2x-1) \\ \Rightarrow 3x < 2\sqrt{3} + 4x - 2\end{aligned}$$

-2-

## 1YGS-MPI PAPER 1 - QUESTION 8

$$\Rightarrow -x < -2 + 2\sqrt{3}$$
$$\Rightarrow \underline{x > 2 - 2\sqrt{3}} \quad \text{MULTIPLY BY NEGATN}$$

FINALLY TO OBTAIN THE COMMON SOLUTION INTERVAL



$$\therefore \underline{2 - 2\sqrt{3} < x < 2 - \sqrt{3}} \quad \text{OR} \quad \underline{x > 2 + \sqrt{3}}$$



-1-


## IYGB - MPI PAPER K - QUESTION 9

OBTAIN THE COORDINATES OF A & B

$$\begin{aligned} y &= 10 - 4x - x^2 \\ y &= 6 - x \end{aligned} \quad \left. \begin{array}{l} \Rightarrow 6 - x = 10 - 4x - x^2 \\ \Rightarrow x^2 + 3x - 4 = 0 \\ \Rightarrow (x+4)(x-1) = 0 \\ \Rightarrow x = -4, 1 \quad y = 10, 5 \end{array} \right\}$$

$$\therefore A(-4, 10) \quad B(1, 5)$$

LOOKING AT THE PICTORIAL EQUATION BELOW



$$\begin{aligned} &\int_{-4}^1 (10 - 4x - x^2) dx \\ &= \left[ 10x - 2x^2 - \frac{1}{3}x^3 \right]_{-4}^1 \end{aligned}$$

$$= (10 - 2 - \frac{1}{3}) - (-40 - 32 + \frac{64}{3})$$

$$= \frac{23}{3} - \left( -\frac{152}{3} \right)$$

$$= \frac{175}{3}$$

$$\frac{1}{2}(10+5) \times 5 = \frac{75}{2}$$

$$\text{REQUIRED AREA} = \frac{175}{3} - \frac{75}{2} = \frac{125}{6}$$

- 1 -

## IYGB-MPI PAPER K - QUESTION 10

SOLVING SIMULTANEOUSLY

$$y = 9^x \quad \& \quad y = 6 \times 5^x$$

$$\downarrow \qquad \downarrow$$

$$9^x = 6 \times 5^x$$

TAKING LOGARITHMS BASE 3

$$\Rightarrow \log_3 9^x = \log_3 (6 \times 5^x)$$

$$\Rightarrow x \log_3 9 = \log_3 6 + \log_3 5^x$$

$$\Rightarrow x \log_3 9 = \log_3 6 + x \log_3 5$$

$$\Rightarrow x \log_3 9 - x \log_3 5 = \log_3 6$$

$$\Rightarrow x [\log_3 9 - \log_3 5] = \log_3 6$$

$$\Rightarrow x = \frac{\log_3 6}{\log_3 9 - \log_3 5}$$

TIDY FURTHER AS FOLLOWS


$$\Rightarrow x = \frac{\log_3 3 + \log_3 2}{\log_3 3^2 - \log_3 5}$$

$$\Rightarrow x = \frac{1 + \log_3 2}{2 \log_3 3 - \log_3 5}$$

$$\Rightarrow x = \frac{1 + \log_3 2}{2 - \log_3 5}$$

IYGB - MPI PAPER K - QUESTION 11

a)



CONSTRAINT SURFACE AREA =  $192\pi$

$$\Rightarrow h + \text{rectangle} + 2\textcircles = 192\pi$$

$$\Rightarrow 2\pi rh + 2 \times \pi r^2 = 192\pi$$

$$\Rightarrow rh + r^2 = 96$$

Volume =  $\pi r^2 \times h$

$$\Rightarrow V = \pi r(rh)$$

$$\Rightarrow V = \pi r(96 - r^2)$$

$$\Rightarrow V = 96\pi r - \pi r^3$$

$$rh = 96 - r^2$$

AS REQUIRED

b)

Differentiate & solve for zero

$$\Rightarrow V = 96\pi r - \pi r^3$$

$$\Rightarrow \frac{dV}{dr} = 96\pi - 3\pi r^2$$

$$\Rightarrow 0 = 96\pi - 3\pi r^2$$

$$\Rightarrow 3\pi r^2 = 96\pi$$

$$\Rightarrow r^2 = 32$$

$$\Rightarrow r = +\sqrt{32} \approx 5.66 \text{ cm}$$

-2-

## LYGB - MPI PAPER K - QUESTION 11

c) CHECKING WITH THE 2ND DERIVATIVE

$$\Rightarrow \frac{dV}{dr} = 96\pi - 3\pi r^2$$

$$\Rightarrow \frac{d^2V}{dr^2} = -6\pi r$$

$$\Rightarrow \left. \frac{d^2V}{dr^2} \right|_{r=5.65} = -106.62... < 0$$

indeed it will give  
the maximum value for V

d)

$$V = 96\pi r - \pi r^3$$


$$\Rightarrow V_{MAX} = 96\pi(4\sqrt{2}) - \pi(4\sqrt{2})^3$$

$$\Rightarrow V_{MAX} = 4\pi\sqrt{2} [96 - (4\sqrt{2})^2]$$

$$\Rightarrow V_{MAX} = 4\pi\sqrt{2} \times 64$$

$$\Rightarrow V_{MAX} = 256\pi\sqrt{2}$$

$$\Rightarrow V_{MAX} \approx 1137$$



-1-

## IYG-B - MPI PAPER K - QUESTION 12

a) PROCEED AS FOLLOWS

- GRADIENT  $AB = \frac{y_2 - y_1}{x_2 - x_1} = \frac{5 - 1}{6 - 0} = \frac{4}{6} = \frac{2}{3}$
- OUR LNE (PERPENDICULAR BISECTOR) =  $-\frac{3}{2}$
- FINDING THE MIDPOINT OF AB :  $\left(\frac{0+6}{2}, \frac{1+5}{2}\right) = (3, 3)$
- REQUIRED LNE HAS EQUATION

$$\begin{aligned} y - y_0 &= m(x - x_0) \\ y - 3 &= -\frac{3}{2}(x - 3) \\ 2y - 6 &= -3x + 9 \\ 2y + 3x &= 15 \end{aligned}$$



//

b) LOOKING AT A DIAGRAM - NOT DRAWN TO SCALE

$$\text{GRADIENT } OA = \frac{y_2 - y_1}{x_2 - x_1} = \frac{1 - 0}{0 - 0} = \frac{1}{0}$$

i.e INFINITE GRADIENT

i.e LNE IS VERTICAL



∴ PERPENDICULAR BISECTOR HAS EQUATION  $y = \frac{1}{2}$

USING  $2y + 3x = 15$  WITH  $y = \frac{1}{2}$

$$2 \times \frac{1}{2} + 3x = 15$$


$$\begin{aligned} 3x &= 14 \\ x &= \frac{14}{3} \end{aligned}$$

$$\therefore C\left(\frac{14}{3}, \frac{1}{2}\right)$$

-1-

## LYGB - MPI PAPER 1 - QUESTION 13

a) LOOKING AT A SKETCH OF THE GRAPH OF  $y = \cos x$



TRANSLATING BY  $60^\circ$  TO THE "LEFT"

$$\therefore (270, 0) \mapsto B(210, 0)$$

$$(360, 1) \mapsto C(300, 0)$$

of SUBSTITUTING  $x=0$  INTO  $y = \cos(x+60)$  GIVES  $y = \frac{1}{2}$

$$\therefore A(0, \frac{1}{2})$$

b) SOLVING THE EQUATION  $y = -\frac{1}{2}$

$$\Rightarrow \cos(x+60) = -\frac{1}{2}$$

$$\arccos\left(-\frac{1}{2}\right) = 120^\circ$$

$$\Rightarrow \begin{cases} x+60^\circ = 120^\circ + 360n \\ x+60^\circ = 240^\circ + 360n \end{cases} \quad n=0, 1, 2, 3, \dots$$

$$\Rightarrow \begin{cases} x = 60^\circ + 360n \\ x = 180^\circ + 360n \end{cases}$$

$$\therefore x_1 = 60^\circ$$

$$x_2 = 180^\circ$$

$$\text{at } P(60^\circ, \frac{1}{2}) \text{ & } Q(180^\circ, \frac{1}{2})$$