

Pure 34 – Compound Angles

Please **complete** this homework by _____. Start it early. If you can't do a question you will then have time to ask your teacher for help or go to a drop in session.

Section 1 – Review of previous topics. Please **complete** all questions.

1. $f(x) = x^2 - 8x + 19$

- (a) Express $f(x)$ in the form $(x + a)^2 + b$, where a and b are constants.

The curve C with equation $y = f(x)$ crosses the y -axis at the point P and has a minimum point at the point Q .

- (b) Sketch the graph of C showing the coordinates of point P and the coordinates of point Q .
 (c) Find the distance PQ , writing your answer as a simplified surd.

2. A circle C has centre $(-1, 7)$ and passes through the point $(0, 0)$. Find an equation for C .

3.




Figure 1

Figure 1 shows a sketch of the curve with equation $y = f(x)$ where

$$f(x) = \frac{x}{x-2}, \quad x \neq 2.$$

The curve passes through the origin and has two asymptotes, with equations $y = 1$ and $x = 2$, as shown in Figure 1.

- (a) Sketch the curve with equation $y = f(x - 1)$ and state the equations of the asymptotes of this curve.
 (b) Find the coordinates of the points where the curve with equation $y = f(x - 1)$ crosses the coordinate axes.

4. Solve for x in the interval $0 \leq x \leq 2\pi$ the following equation, giving your answer in terms of π

$$\operatorname{cosec}^2 x + \cot^2 x = 3$$

5. Prove the following identities
- $\sec^2 x - \sin^2 x \equiv \tan^2 x + \cos^2 x$
 - $(\sin x - \sec x)^2 \equiv \sin^2 x + (\tan x - 1)^2$

Section 2 – Consolidation of this week’s topic. Please complete all questions.

Total 51

- Express in the form $\sin \alpha$, where α is acute.
 - $\sin 10^\circ \cos 30^\circ + \cos 10^\circ \sin 30^\circ$
 - $\cos 14^\circ \cos 39^\circ - \sin 14^\circ \sin 39^\circ$ **(3)**
- Express as a single trigonometric ratio

$$\frac{\tan 2A + \tan 5}{1 - \tan 2A \tan 5A}$$
 (1)
- Find the maximum value that each expression can take and the smallest positive value of x , in degrees for which this maximum occurs
 - $\cos x \cos 30^\circ + \sin x \sin 30^\circ$
 - $3 \sin x \cos 45^\circ + 3 \cos x \sin 45^\circ$ **(6)**
- Find the minimum value that each expression can take and the smallest positive value of x , in radians in terms of π , for which this minimum occurs.
 - $\sin x \cos \frac{\pi}{3} - \cos x \sin \frac{\pi}{3}$
 - $2 \cos x \cos \frac{\pi}{6} - 2 \sin x \sin \frac{\pi}{6}$ **(6)**
- Solve each equation for θ in the interval $0 \leq \theta \leq 360$.
Give your answers to 1 decimal place where appropriate.
 - $\sin \theta \cos 15^\circ + \cos \theta \sin 15^\circ = 0.4$ **(3)**
 - $\frac{\tan 2\theta - \tan 6}{1 + \tan 2\theta \tan 6} = 1$ **(5)**
 - $\cos(\theta - 60) = \sin \theta$ **(6)**
- Given that

$$2 \cos(x + 50)^\circ = \sin(x + 40)^\circ.$$
 - Show that

$$\tan x^\circ = \frac{1}{3} \tan 40^\circ.$$
 (4)
 - Hence solve, for $0 \leq \vartheta < 360$,

$$2 \cos(2\vartheta + 50)^\circ = \sin(2\vartheta + 40)^\circ,$$
 giving your answers to 1 decimal place. **(4)**

- 7) (a) Starting from the formulae for $\sin(A + B)$ and $\cos(A + B)$, prove that

$$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}.$$

(4)

- (b) Deduce that

$$\tan\left(\theta + \frac{\pi}{6}\right) = \frac{1 + \sqrt{3} \tan \theta}{\sqrt{3} - \tan \theta}.$$

(3)

- (c) Hence, or otherwise, solve, for $0 \leq \vartheta \leq \pi$,
 $1 + \sqrt{3} \tan \vartheta = (\sqrt{3} - \tan \vartheta) \tan(\pi - \vartheta)$.
Give your answers as multiples of π .

(6)