

Surname		Other Names	
Centre Number		Candidate Number	
Candidate Signature			

Leave blank
-------------

General Certificate of Education  
January 2005  
Advanced Level Examination



**PHYSICS (SPECIFICATION A) PHA8/W**  
**Unit 8 Nuclear Instability: Turning Points in Physics Option**

Wednesday 26 January 2005 Morning Session

**In addition to this paper you will require:**

- a calculator;
- a pencil and a ruler.

For Examiner's Use			
Number	Mark	Number	Mark
1			
2			
3			
4			
5			
Total (Column 1)	→		
Total (Column 2)	→		
TOTAL			
Examiner's Initials			

Time allowed: 1 hour 15 minutes

**Instructions**

- Use blue or black ink or ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions in the spaces provided. All working must be shown.
- Do all rough work in this book. Cross through any work you do not want marked.

**Information**

- The maximum mark for this paper is 40.
- Mark allocations are shown in brackets.
- The paper carries 10% of the total marks for Physics Advanced.
- A *Data Sheet* is provided on pages 3 and 4. You may wish to detach this perforated sheet at the start of the examination.
- You are expected to use a calculator where appropriate.
- In questions requiring description and explanation you will be assessed on your ability to use an appropriate form and style of writing, to organise relevant information clearly and coherently, and to use specialist vocabulary where appropriate. The degree of legibility of your handwriting and the level of accuracy of your spelling, punctuation and grammar will also be taken into account.

**Data Sheet**

- A perforated *Data Sheet* is provided as pages 3 and 4 of this question paper.
- This sheet may be useful for answering some of the questions in the examination.
- You may wish to detach this sheet before you begin work.

Fundamental constants and values				Mechanics and Applied Physics		Fields, Waves, Quantum Phenomena		
Quantity	Symbol	Value	Units					
speed of light in vacuo	$c$	$3.00 \times 10^8$	$\text{m s}^{-1}$	$v = u + at$	$g = \frac{F}{m}$			
permeability of free space	$\mu_0$	$4\pi \times 10^{-7}$	$\text{H m}^{-1}$	$s = \left(\frac{u+v}{2}\right)t$	$g = -\frac{GM}{r^2}$			
permittivity of free space	$\epsilon_0$	$8.85 \times 10^{-12}$	$\text{F m}^{-1}$	$s = ut + \frac{at^2}{2}$	$g = -\frac{\Delta V}{\Delta x}$			
charge of electron	$e$	$1.60 \times 10^{-19}$	C	$v^2 = u^2 + 2as$	$V = -\frac{GM}{r}$			
the Planck constant	$h$	$6.63 \times 10^{-34}$	J s	$F = \frac{\Delta(mv)}{\Delta t}$	$a = -(2\pi f)^2 x$			
gravitational constant	$G$	$6.67 \times 10^{-11}$	$\text{N m}^2 \text{kg}^{-2}$	$P = Fv$	$v = \pm 2\pi f \sqrt{A^2 - x^2}$			
the Avogadro constant	$N_A$	$6.02 \times 10^{23}$	$\text{mol}^{-1}$	$\text{efficiency} = \frac{\text{power output}}{\text{power input}}$	$x = A \cos 2\pi ft$			
molar gas constant	$R$	8.31	$\text{J K}^{-1} \text{mol}^{-1}$	$\omega = \frac{v}{r} = 2\pi f$	$T = 2\pi \sqrt{\frac{m}{k}}$			
the Boltzmann constant	$k$	$1.38 \times 10^{-23}$	$\text{J K}^{-1}$	$a = \frac{v^2}{r} = r\omega^2$	$T = 2\pi \sqrt{\frac{l}{g}}$			
the Stefan constant	$\sigma$	$5.67 \times 10^{-8}$	$\text{W m}^{-2} \text{K}^{-4}$	$I = \sum mr^2$	$\lambda = \frac{\omega s}{D}$			
the Wien constant	$\alpha$	$2.90 \times 10^{-3}$	m K	$E_k = \frac{1}{2} I\omega^2$	$d \sin \theta = n\lambda$			
electron rest mass	$m_e$	$9.11 \times 10^{-31}$	kg	$\omega_2 = \omega_1 + at$	$\theta \approx \frac{\lambda}{D}$			
(equivalent to $5.5 \times 10^{-4}u$ )				$\theta = \omega_1 t + \frac{1}{2} at^2$	$1/n_2 = \frac{\sin \theta_1}{\sin \theta_2} = \frac{c_1}{c_2}$			
electron charge/mass ratio	$e/m_e$	$1.76 \times 10^{11}$	$\text{C kg}^{-1}$	$\omega_2^2 = \omega_1^2 + 2a\theta$	$1/n_2 = \frac{n_2}{n_1}$			
proton rest mass	$m_p$	$1.67 \times 10^{-27}$	kg	$\theta = \frac{1}{2}(\omega_1 + \omega_2)t$	$\sin \theta_c = \frac{1}{n}$			
(equivalent to 1.00728u)				$T = I\alpha$	$E = hf$			
proton charge/mass ratio	$e/m_p$	$9.58 \times 10^7$	$\text{C kg}^{-1}$	$\text{angular momentum} = I\omega$	$hf = \phi + E_k$			
neutron rest mass	$m_n$	$1.67 \times 10^{-27}$	kg	$W = T\theta$	$hf = E_1 - E_2$			
(equivalent to 1.00867u)				$P = T\omega$	$\lambda = \frac{h}{p} = \frac{h}{mv}$			
gravitational field strength	$g$	9.81	$\text{N kg}^{-1}$	$\text{angular impulse} = \text{change of angular momentum} = Tt$	$c = \frac{1}{\sqrt{\mu_0 \epsilon_0}}$			
acceleration due to gravity	$g$	9.81	$\text{m s}^{-2}$	$\Delta Q = \Delta U + \Delta W$	<b>Electricity</b>			
atomic mass unit	$u$	$1.661 \times 10^{-27}$	kg	$\Delta W = p\Delta V$	$\epsilon = \frac{E}{Q}$			
(1u is equivalent to 931.3 MeV)				$pV^\gamma = \text{constant}$	$\epsilon = I(R+r)$			
<b>Fundamental particles</b>				$\text{work done per cycle} = \text{area of loop}$	$\frac{1}{R_T} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots$			
<i>Class</i>	<i>Name</i>	<i>Symbol</i>	<i>Rest energy</i>	$\text{input power} = \text{calorific value} \times \text{fuel flow rate}$	$R_T = R_1 + R_2 + R_3 + \dots$			
			/MeV	$\text{indicated power as (area of } p-V \text{ loop)} \times (\text{no. of cycles/s}) \times (\text{no. of cylinders})$	$P = I^2 R$			
photon	photon	$\gamma$	0	$\text{friction power} = \text{indicated power} - \text{brake power}$	$E = \frac{F}{Q} = \frac{V}{d}$			
lepton	neutrino	$\nu_e$	0	$\text{efficiency} = \frac{W}{Q_{in}} = \frac{Q_{in} - Q_{out}}{Q_{in}}$	$E = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}$			
		$\nu_\mu$	0	$\text{maximum possible efficiency} = \frac{T_H - T_C}{T_H}$	$E = \frac{1}{2} QV$			
mesons	electron	$e^\pm$	0.510999		$F = BI$			
		muon	$\mu^\pm$	105.659		$F = BQv$		
	pion	$\pi^\pm$	139.576			$Q = Q_0 e^{-t/RC}$		
		$\pi^0$	134.972			$\Phi = BA$		
kaon	$K^\pm$	493.821						
	$K^0$	497.762						
baryons	proton	p	938.257					
	neutron	n	939.551					
<b>Properties of quarks</b>								
<i>Type</i>	<i>Charge</i>	<i>Baryon number</i>	<i>Strangeness</i>					
u	$+\frac{2}{3}$	$+\frac{1}{3}$	0					
d	$-\frac{1}{3}$	$+\frac{1}{3}$	0					
s	$-\frac{1}{3}$	$+\frac{1}{3}$	-1					
<b>Geometrical equations</b>								
arc length = $r\theta$								
circumference of circle = $2\pi r$								
area of circle = $\pi r^2$								
area of cylinder = $2\pi rh$								
volume of cylinder = $\pi r^2 h$								
area of sphere = $4\pi r^2$								
volume of sphere = $\frac{4}{3}\pi r^3$								

$$\text{magnitude of induced e.m.f.} = N \frac{\Delta\Phi}{\Delta t}$$

$$I_{\text{rms}} = \frac{I_0}{\sqrt{2}}$$

$$V_{\text{rms}} = \frac{V_0}{\sqrt{2}}$$

### Mechanical and Thermal Properties

$$\text{the Young modulus} = \frac{\text{tensile stress}}{\text{tensile strain}} = \frac{F}{A} \frac{l}{e}$$

$$\text{energy stored} = \frac{1}{2} Fe$$

$$\Delta Q = mc \Delta\theta$$

$$\Delta Q = ml$$

$$pV = \frac{1}{3} Nmc^2$$

$$\frac{1}{2} mc^2 = \frac{3}{2} kT = \frac{3RT}{2N_A}$$

### Nuclear Physics and Turning Points in Physics

$$\text{force} = \frac{eV_p}{d}$$

$$\text{force} = Bev$$

$$\text{radius of curvature} = \frac{mv}{Be}$$

$$\frac{eV}{d} = mg$$

$$\text{work done} = eV$$

$$F = 6\pi\eta rv$$

$$I = k \frac{I_0}{x^2}$$

$$\frac{\Delta N}{\Delta t} = -\lambda N$$

$$\lambda = \frac{h}{\sqrt{2meV}}$$

$$N = N_0 e^{-\lambda t}$$

$$T_{\frac{1}{2}} = \frac{\ln 2}{\lambda}$$

$$R = r_0 A^{\frac{1}{3}}$$

$$E = mc^2 = \frac{m_0 c^2}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}}$$

$$l = l_0 \left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}$$

$$t = \frac{t_0}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}}$$

### Astrophysics and Medical Physics

Body	Mass/kg	Mean radius/m
Sun	$2.00 \times 10^{30}$	$7.00 \times 10^8$
Earth	$6.00 \times 10^{24}$	$6.40 \times 10^6$

Sun	$2.00 \times 10^{30}$	$7.00 \times 10^8$
Earth	$6.00 \times 10^{24}$	$6.40 \times 10^6$

1 astronomical unit =  $1.50 \times 10^{11}$  m

1 parsec = 206265 AU =  $3.08 \times 10^{16}$  m = 3.26 ly

1 light year =  $9.45 \times 10^{15}$  m

Hubble constant ( $H$ ) =  $65 \text{ km s}^{-1} \text{ Mpc}^{-1}$

$$M = \frac{\text{angle subtended by image at eye}}{\text{angle subtended by object at unaided eye}}$$

$$M = \frac{f_o}{f_c}$$

$$m - M = 5 \log \frac{d}{10}$$

$$\lambda_{\text{max}} T = \text{constant} = 0.0029 \text{ m K}$$

$$v = Hd$$

$$P = \sigma AT^4$$

$$\frac{\Delta f}{f} = \frac{v}{c}$$

$$\frac{\Delta \lambda}{\lambda} = -\frac{v}{c}$$

$$R_s \approx \frac{2GM}{c^2}$$

### Medical Physics

$$\text{power} = \frac{1}{f}$$

$$\frac{1}{u} + \frac{1}{v} = \frac{1}{f} \text{ and } m = \frac{v}{u}$$

$$\text{intensity level} = 10 \log \frac{I}{I_0}$$

$$I = I_0 e^{-\mu x}$$

$$\mu_m = \frac{\mu}{\rho}$$

### Electronics

Resistors

Preferred values for resistors (E24)  
Series: 1.0 1.1 1.2 1.3 1.5 1.6 1.8 2.0 2.2 2.4 2.7 3.0 3.3 3.6 3.9 4.3 4.7 5.1 5.6 6.2 6.8 7.5 8.2 9.1 ohms  
and multiples that are ten times greater

$$Z = \frac{V_{\text{rms}}}{I_{\text{rms}}}$$

$$\frac{1}{C_T} = \frac{1}{C_1} + \frac{1}{C_2} + \frac{1}{C_3} + \dots$$

$$C_T = C_1 + C_2 + C_3 + \dots$$

$$X_C = \frac{1}{2\pi f C}$$

### Alternating Currents

$$f = \frac{1}{T}$$

### Operational amplifier

$$G = \frac{V_{\text{out}}}{V_{\text{in}}} \quad \text{voltage gain}$$

$$G = -\frac{R_f}{R_i} \quad \text{inverting}$$

$$G = 1 + \frac{R_f}{R_i} \quad \text{non-inverting}$$

$$V_{\text{out}} = -R_f \left( \frac{V_1}{R_1} + \frac{V_2}{R_2} + \frac{V_3}{R_3} \right) \quad \text{summing}$$

## SECTION A: NUCLEAR INSTABILITY

Answer **all** of this question

- 1 The high energy electron diffraction apparatus represented in **Figure 1** can be used to determine nuclear radii. The intensity of the electron beam received by the detector is measured at various diffraction angles,  $\theta$ .




Figure 1

- (a) Sketch on the axes below a graph of the results expected from such an electron diffraction experiment.



Turn over ►

- (b) (i) Use the data in the table to plot a straight line graph that confirms the relationship  $R = r_0 A^{\frac{1}{3}}$ .

element	radius of nucleus, $R$ $10^{-15}\text{m}$	nucleon number, $A$	
lead	6.66	208	
tin	5.49	120	
iron	4.35	56	
silicon	3.43	28	
carbon	2.66	12	

- (ii) Estimate the value of  $r_0$  from the graph.

.....

.....

.....

.....

.....

(5 marks)

- (c) Discuss the merits of using high energy electrons to determine nuclear radii rather than using  $\alpha$  particles.

You may be awarded marks for the quality of written communication in your answer.

.....

.....


.....

.....

.....

.....

(3 marks)




Turn over ►

**SECTION B: TURNING POINTS IN PHYSICS**

Answer **all** questions.

2 A narrow beam of electrons is produced in a vacuum tube using the arrangement shown in **Figure 2**.



**Figure 2**

(a) Describe the function of each voltage supply unit and state a typical voltage for each unit.

(i) unit A

.....

.....

(ii) unit B

.....

.....

*(3 marks)*

(b) State and explain the effect on the beam of

(i) reducing the voltage of A,

.....

.....

.....

.....

.....



(ii) increasing the voltage of B.

.....

.....

.....

.....

.....

(4 marks)

7

**TURN OVER FOR THE NEXT QUESTION**

**Turn over ▶**

- 3 In an experiment to demonstrate the wave nature of light, a parallel beam of monochromatic light was directed at two closely spaced slits, as shown in **Figure 3**. A pattern of bright and dark fringes due to this light passing through the slits was seen on the screen.



**Figure 3**

- (a) Explain why this fringe pattern was formed.

You may be awarded marks for the quality of written communication in your answer.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(4 marks)

(b) Discuss why this fringe pattern cannot be explained using Newton's corpuscular theory of light.

.....

.....

.....

.....

.....


(2 marks)

6

**TURN OVER FOR THE NEXT QUESTION**

**Turn over ▶**

- 4 A charged oil droplet was observed falling between two oppositely charged parallel plates, as shown in **Figure 4**.



**Figure 4**

- (a) Explain why the droplet stopped moving and remained stationary when the potential difference between the plates was adjusted to a certain value,  $V_c$ .

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(3 marks)

- (b) (i) The spacing between the plates is 6.0 mm. A charged oil droplet of mass  $6.2 \times 10^{-14}$  kg is stopped when  $V_c = 5700$  V. Calculate the charge on this droplet.

.....

.....

.....

.....

.....

- (ii) Describe and explain what would have happened to this droplet if the potential difference had been greater than 5700 V.

.....

.....

.....

.....

.....

(5 marks)

8

**TURN OVER FOR THE NEXT QUESTION**

**Turn over ▶**

5 (a) In a science fiction film, a space rocket travels away from the Earth at a speed of  $0.994c$ , where  $c$  is the speed of light in free space. A radio message of duration 800 s is transmitted by the space rocket.

(i) Calculate the duration of the message when it is received at the Earth.

.....  
.....  
.....  
.....

(ii) Calculate the distance moved by the rocket in the Earth's frame of reference in the time taken to send the message.

.....  
.....  
.....

(4 marks)

(b) A student claims that a twin who travels at a speed close to the speed of light from Earth to a distant star and back would, on return to Earth, be a different age to the twin who stayed on Earth. Discuss whether or not this claim is correct.

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

(3 marks)

**QUALITY OF WRITTEN COMMUNICATION**

**END OF QUESTIONS**

7

2