

1 $= 1 + 4(4x) + 6(4x)^2 + 4(4x)^3 + (4x)^4$
 $= 1 + 16x + 96x^2 + 256x^3 + 256x^4$

2 a $u_5 = 3 \times (-2)^4 = 48$
b $S_{10} = \frac{3[1 - (-2)^{10}]}{1 - (-2)} = -1023$
c positive terms form GP:
 $a = 3, r = (-2)^2 = 4$
 $S_8 = \frac{3(4^8 - 1)}{4 - 1} = 65\ 535$

3 a $= 1 + 7(3x) + \frac{7 \times 6}{2} (3x)^2$
 $+ \frac{7 \times 6 \times 5}{3 \times 2} (3x)^3 + \frac{7 \times 6 \times 5 \times 4}{4 \times 3 \times 2} (3x)^4 + \dots$
 $= 1 + 21x + 189x^2 + 945x^3 + 2835x^4 + \dots$
b let $x = 0.01$
 $1.03^7 \approx 1 + 0.21 + 0.0189$
 $+ 0.000\ 945 + 0.000\ 028\ 35$
 $= 1.229\ 87$ (5dp)

4 GP: $a = 8, r = 2, n = 10$
 $S_{10} = \frac{8(2^{10} - 1)}{2 - 1} = 8184$

5 a $= 2^5 + 5(2^4)x + 10(2^3)x^2$
 $+ 10(2^2)x^3 + 5(2)x^4 + x^5$
 $= 32 + 80x + 80x^2 + 40x^3 + 10x^4 + x^5$
b $= 32 - 80x + 80x^2 - 40x^3 + 10x^4 - x^5$
c $(2 + \sqrt{5})^5 = 32 + 80(\sqrt{5}) + 80(\sqrt{5})^2$
 $+ 40(\sqrt{5})^3 + 10(\sqrt{5})^4 + (\sqrt{5})^5$
 $= 32 + 80\sqrt{5} + 400 + 200\sqrt{5} + 250 + 25\sqrt{5}$
 $= 682 + 305\sqrt{5}$
 $\therefore (2 + \sqrt{5})^5 - (2 - \sqrt{5})^5$
 $= (682 + 305\sqrt{5}) - (682 - 305\sqrt{5})$
 $= 610\sqrt{5}, k = 610$

6 a amount in account after 3rd payment in
 $= 200 + (1.005 \times 200) + (1.005^2 \times 200)$
 $= 603.005$
interest paid at end of 3rd month
 $= 0.005 \times 603.005 = £3.02$ (nearest penny)
b amount paid in $= 12 \times 200 = £2400$
amount in account after 12 months
 $= 200(1.005 + 1.005^2 + \dots + 1.005^{12})$
 $= 200 \times S_{12}$ [GP: $a = 1.005, r = 1.005$]
 $= 200 \times \frac{1.005(1.005^{12} - 1)}{1.005 - 1} = 2479.45$
total interest $= 2479.45 - 2400 = £79.45$

7 $= 1 + 8(-3x) + \frac{8 \times 7}{2} (-3x)^2$
 $+ \frac{8 \times 7 \times 6}{3 \times 2} (-3x)^3 + \dots$
 $= 1 - 24x + 252x^2 - 1512x^3 + \dots$

8 a $S_n = a + ar + ar^2 + \dots + ar^{n-1}$
 $rS_n = ar + ar^2 + \dots + ar^{n-1} + ar^n$
subtracting, $S_n - rS_n = a - ar^n$
 $S_n(1 - r) = a(1 - r^n)$
 $S_n = \frac{a(1 - r^n)}{1 - r}$

b $r = 6 \div 3 = 2$
 $a \times 2^3 = 3 \therefore a = \frac{3}{8}$
 $S_{16} = \frac{\frac{3}{8}(2^{16} - 1)}{2 - 1} = 24\ 575\ \frac{5}{8}$

9 **a** $= 1 + n(ax) + \frac{n(n-1)}{2}(ax)^2 + \dots$
 $= 1 + anx + \frac{1}{2}a^2n(n-1)x^2 + \dots$

b $\frac{1}{2}a^2n(n-1) = 3an$
 $a^2n(n-1) = 6an$
 $an[a(n-1) - 6] = 0$
 $n \neq 0 \therefore a(n-1) - 6 = 0$
 $an - a = 6$
 $n = \frac{6+a}{a}$

c $n = 10 \therefore \text{coeff. of } x^3 = \frac{10 \times 9 \times 8}{3 \times 2} \times (\frac{2}{3})^3 = 35\frac{5}{9}$

11 **a** $\frac{162}{1-r} = 486$
 $1-r = \frac{162}{486} = \frac{1}{3} \therefore r = \frac{2}{3}$

b $u_6 = 162 \times (\frac{2}{3})^5 = \frac{64}{3}$ or $21\frac{1}{3}$

c $S_{10} = \frac{162[1-(\frac{2}{3})^{10}]}{1-\frac{2}{3}} = 477.572$

13 **a** time $= 120 \times (0.9)^3 = 87.48$ seconds

b GP: $a = 120, r = 0.9, n = 12$
 $S_{12} = \frac{120[1-(0.9)^{12}]}{1-0.9}$
 $= 861.08$ seconds
 $= 14$ mins 21 secs (nearest sec.)

15 **a** 6, 12, 24, 48

b GP: $a = 6, r = 2, n = 10$
 $S_{10} = \frac{6(2^{10}-1)}{2-1} = 6138$

17 **a** $a \times (1.5)^2 = 18$
 $a = 18 \div 2.25 = 8$

b $S_6 = \frac{8[(1.5)^6 - 1]}{1.5 - 1} = 166.25$

c $8 \times (1.5)^{k-1} > 8000$
 $(k-1) \lg 1.5 > \lg 1000$
 $k > \frac{\lg 1000}{\lg 1.5} + 1$
 $k > 18.04 \therefore \text{smallest } k = 19$

10 $= 2^6 + 6(2^5)(5x) + \frac{6 \times 5}{2}(2^4)(5x)^2 + \dots$
 $= 64 + 960x + 6000x^2 + \dots$

12 **a** $= 1 + 4(3x) + 6(3x)^2 + 4(3x)^3 + (3x)^4$
 $= 1 + 12x + 54x^2 + 108x^3 + 81x^4$

b term in $x^2 = (1)(54x^2) + (4x)(12x) + (-x^2)(1)$
coefficient of $x^2 = 54 + 48 - 1 = 101$

14 $= [1+8(\frac{x}{2})+\frac{8 \times 7}{2}(\frac{x}{2})^2+\dots][1+6(-x)+\frac{6 \times 5}{2}(-x)^2+\dots]$
 $= [1+4x+7x^2+\dots][1-6x+15x^2+\dots]$
 $= 1-6x+15x^2+4x-24x^2+7x^2+\dots$
 $= 1-2x-2x^2+\dots$
 $\therefore A = -2, B = -2$

16 **a** $= 1 + 4x + 6x^2 + 4x^3 + x^4$
b $= 1 - 4x + 6x^2 - 4x^3 + x^4$
c $(1 + 4x + 6x^2 + 4x^3 + x^4)$
 $+ (1 - 4x + 6x^2 - 4x^3 + x^4) = 82$
 $2 + 12x^2 + 2x^4 = 82$
 $x^4 + 6x^2 - 40 = 0$
 $(x^2 + 10)(x^2 - 4) = 0$
 $x^2 = -10$ [no real solutions] or $x^2 = 4$
 $x = \pm 2$

18 $(1 + \frac{ax}{2})^{10} + (1 + bx)^{10}$
 $= 1 + 10(\frac{ax}{2}) + \frac{10 \times 9}{2}(\frac{ax}{2})^2 + \dots$
 $+ 1 + 10(bx) + \frac{10 \times 9}{2}(bx)^2 + \dots$
 $= 2 + (5a + 10b)x + (\frac{45}{4}a^2 + 45b^2)x^2 + \dots$
 $\therefore 5a + 10b = 0 \Rightarrow a = -2b$
and $\frac{45}{4}a^2 + 45b^2 = 90 \Rightarrow a^2 + 4b^2 = 8$
sub. $(-2b)^2 + 4b^2 = 8$
 $b^2 = 1$
 $a < b \therefore b = 1, a = -2$